INTERNATIONAL JOURNAL OF CLINICAL, BIOMEDICAL AND HEALTH SCIENCES RESEARCH
2026, Vol 1 Issue 1, pp. 21-24
ISSN:*xxxx-xxxx*
Open Access. Online

www.ijcbhsr.com/

Lifestyle Factors and Cardiovascular Risk

Dr. Sophia Claire Robinson

Assistant Professor, School of Financial Technology and Innovation,
Western Pacific University, Perth, Australia.

International Journal of Clinical, Biomedical and Health Sciences Research

vol 1, Issue1, (2026).

Received: Jan 24, 2026.

Accepted: April 28, 2026.

Published: June 4, 2026.

ABSTRACT:

Cardiovascular risk reflects an interaction among behavior, biological measures, inherited susceptibility and the social environment. This narrative review examines tobacco exposure, diet, physical activity, sleep, alcohol use and psychosocial conditions, and explains how they relate to blood pressure, lipids, glucose and body weight. It synthesizes World Health Organization guidance, the American Heart Association's Life's Essential 8 framework and the multicountry INTERHEART case-control study. A chart reproduces selected adjusted odds ratios from INTERHEART, with their distinct comparison groups stated; these associations are neither individualized risk estimates nor proof of causal effects. Tables summarize pathways and practical assessment measures. Evidence supports a combined approach that helps people avoid tobacco, eat a healthier dietary pattern, move regularly and maintain healthy sleep while monitoring clinical risk factors. Individual choices depend on access, income, work, environment and health status; prevention therefore also requires supportive systems and appropriate clinical care [1–5].

Keywords: cardiovascular disease; lifestyle; smoking; dietary pattern; physical activity; sleep; prevention; risk factors.

1 Introduction

Cardiovascular diseases include coronary heart disease and stroke, among other disorders of the heart and blood vessels. The World Health Organization identifies unhealthy diet, physical inactivity, tobacco use and harmful alcohol use as major behavioral risk factors. Their effects may appear as elevated blood pressure, blood glucose and blood lipids, as well as overweight or obesity [1]. These measured factors help clinicians assess risk, but a person's history, age and other conditions also matter. This article examines how lifestyle-related exposures fit into that larger picture.

A useful framework must separate behavior from a health measure. Eating patterns and activity are behaviors; LDL cholesterol and blood pressure are intermediate physiological factors. Body mass index

© The Author(s) 2026. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

reflects a body-size measure rather than a specific behavior. The American Heart Association's Life's Essential 8 includes four health behaviors—diet, physical activity, nicotine exposure and sleep—and four health factors—body mass index, blood lipids, blood glucose and blood pressure [2]. The framework describes cardiovascular health across the life course; it is not a substitute for clinical diagnosis or an individualized treatment plan.

2 Review approach and interpretation

This is a focused narrative review of selected primary studies and authoritative guidance, not a new cohort analysis or meta-analysis. The strongest quantitative illustration is INTERHEART, a case-control study of 15,152 acute myocardial infarction cases and 14,820 controls in 52 countries [3]. Case-control odds ratios quantify associations under specified exposure definitions. They should not be read as the amount by which one individual's future risk will change if a behavior changes. Reverse causation, measurement differences, residual confounding and different baseline risks limit direct translation.

Population attributable risk estimates, when reported, depend on exposure prevalence as well as the strength of an association. They should not be added across correlated exposures. Lifestyle patterns cluster: access to affordable food, safe places to exercise, reliable sleep and smoking cessation support can all vary together. This context is relevant to both research design and public-health implementation.

3 Tobacco and nicotine exposure

Smoking increases cardiovascular risk through vascular injury, inflammation, thrombosis and adverse effects on blood pressure and lipid metabolism. INTERHEART reported an adjusted odds ratio of 2.87 for current compared with never smoking in its acute myocardial infarction analysis [3]. This is a study-level association for that particular comparison, not an absolute probability. Avoiding tobacco initiation and supporting cessation are central prevention strategies. Exposure to secondhand smoke also matters; a comprehensive risk assessment should ask about nicotine and smoke exposure rather than only daily cigarette count.

Cessation often requires more than information. Access to evidence-based behavioral support, appropriate medicines and smoke-free environments can help. Patients with existing cardiovascular disease should also receive clinical guidance about secondary prevention. New nicotine products require careful interpretation because absence of combustion does not mean absence of nicotine exposure or established long-term cardiovascular safety. The article does not treat a product switch as equivalent to quitting.

4 Diet, blood pressure and lipids

Dietary patterns affect cardiovascular health through sodium intake, blood pressure, blood lipids, glucose regulation and energy balance. AHA dietary guidance emphasizes overall patterns rather than isolated nutrients: vegetables and fruits, whole grains, healthy protein sources, unsaturated oils and less heavily processed food can form a sustainable pattern [4]. The exact pattern must fit cultural preferences and food access. Replacing a poor-quality diet with a named diet label alone has little value if the overall nutrient pattern does not improve.

Reducing excess sodium is particularly relevant to elevated blood pressure. Managing saturated fat intake and favoring healthier fat sources may improve lipid profiles. These measures complement, rather than replace, medicines when pharmacological treatment is indicated. In INTERHEART, daily fruit and vegetable consumption was associated with lower myocardial infarction odds (OR 0.70 for daily versus less frequent intake), but this observational estimate is not the causal effect of adding one serving of produce [3].

Table 1. Broad pathways and assessment examples. These are not diagnostic criteria or individual risk estimates.

Factor	Potential pathway	Useful assessment
Tobacco exposure	Vascular injury and thrombosis	Smoking status and exposure history
Dietary pattern	Blood pressure, lipids, glucose	Pattern and sodium-rich foods
Physical inactivity	Fitness, metabolism, blood pressure	Weekly activity and sitting time
Sleep	Physiological regulation and behaviors	Duration, regularity, symptoms
Harmful alcohol use	Blood pressure and other harms	Amount, pattern and context

5 Physical activity and sedentary time

Regular movement supports cardiorespiratory fitness and helps control blood pressure, glucose and body weight. WHO guidelines recommend adults undertake 150–300 minutes of moderate-intensity aerobic activity, or 75–150 minutes of vigorous-intensity activity, or an equivalent combination each week; muscle-strengthening activity is also recommended [5]. Some activity is beneficial when the full target is not yet feasible. An individual starting after illness or with symptoms should select an appropriate plan with a clinician.

In INTERHEART, regular physical activity was associated with lower odds of acute myocardial infarction (OR 0.86 under the study's definition) [3]. That estimate cannot be compared directly with the smoking odds ratio to rank intervention priorities: exposure contrasts and baseline risks differ. Physical activity is also influenced by neighborhood safety, work demands, mobility and access to recreation. Programs that change environments can make sustained behavior easier than relying only on motivation.

6 Sleep, alcohol and psychosocial conditions

Sleep is included in Life's Essential 8 as a distinct component of cardiovascular health [2]. Short, irregular or poor-quality sleep can coexist with stress, shift work and sleep disorders. A sleep assessment should look beyond a single hours-per-night number and consider symptoms such as persistent daytime sleepiness or possible sleep apnea. The response may require changes in schedules, behavior or clinical evaluation; a single recommendation will not address every cause.

WHO lists harmful alcohol use among important behavioral risks [1]. INTERHEART reported an association for regular alcohol consumption in its particular study population, but that result should not be presented as advice to start drinking: observational comparisons are vulnerable to confounding and do not establish a safe cardiovascular dose for every person [3]. Reducing harmful use is the appropriate public-health message. Psychosocial factors were also associated with myocardial infarction in INTERHEART (OR 2.67), but its combined measure should not be equated with an individual's brief episode of stress [3]. Workplace conditions, financial strain and social support can affect both behavior and health.

7 Selected findings from INTERHEART




Figure 1. Selected adjusted odds ratios reported in INTERHEART for acute myocardial infarction [3]. Current versus never smoking, psychosocial factors, history of diabetes or hypertension, regular activity and daily fruit/vegetable intake use different exposure comparisons. The dashed line is OR = 1. Values are associations from a case-control study, not causal treatment effects or comparable individual risk predictions.

Table 2. Selected values reproduced from the INTERHEART abstract [3]; odds ratios are not presented as absolute risks.

Study comparison	Adjusted OR	Interpretive limit
Current vs never smoking	2.87	Exposure history differs by person
Psychosocial factors	2.67	Composite study definition
History of diabetes	2.37	Clinical history, not lifestyle alone
History of hypertension	1.91	Measured status and treatment vary
Regular activity	0.86	Study-specific activity definition
Daily fruit and vegetables	0.70	Dietary pattern may confound

The chart is intentionally selective. INTERHEART also analyzed lipid ratios, abdominal obesity and other exposures. Its reported associations were observed across multiple world regions, yet the case-control design and historical period mean that modern local risk assessment should rely on current clinical measurements and validated tools when available. Risk prediction for a patient requires age, blood pressure, lipid values, diabetes and smoking history, among other inputs, and is separate from this descriptive chart.

8 Prevention in clinical and community settings

A practical prevention pathway combines behavior support with repeated measurement. In primary care, clinicians can assess nicotine exposure, diet, activity and sleep while measuring blood pressure and, when indicated, lipids and glucose. Shared decisions should account for an individual's preferences, medical conditions and ability to act on recommendations. Lifestyle measures and medication are often complementary: a person at high risk should not defer indicated treatment while waiting for a behavioral change to alter a laboratory result.

Public policy determines whether healthy choices are feasible. Affordable nutritious food, smoke-free spaces, safe walking routes and working conditions that permit adequate sleep create opportunities to maintain behavior change. WHO notes the role of supportive environments and air-quality improvement in cardiovascular prevention [1]. Equity therefore matters when comparing communities or designing interventions. A program that assumes abundant time, safe streets and easy access to fresh food may widen rather than narrow health differences.

9 Conclusion

Lifestyle factors influence cardiovascular risk through overlapping biological and social pathways. Tobacco avoidance, a healthier dietary pattern, regular activity and healthy sleep belong in a comprehensive prevention strategy, while harmful alcohol use should be addressed. Blood pressure, glucose, lipids and body weight provide additional measurable information that can guide care. Evidence from INTERHEART shows important associations across countries, and Life's Essential 8 offers a coherent framework for monitoring cardiovascular health [2,3]. Neither source makes an individual's outcome predictable from one behavior alone. Effective prevention combines sustained behavior support, clinical assessment and treatment when indicated, and environments that allow people to make healthier choices.

References

[1] World Health Organization. (2025). Cardiovascular diseases (CVDs). https://www.who.int/news-room/fact-sheets/detail/cardiovascular-diseases-(cvds)

[2] Lloyd-Jones, D. M., et al. (2022). Life's Essential 8: Updating and Enhancing the American Heart Association's Construct of Cardiovascular Health. Circulation, 146(5), e18–e43. https://doi.org/10.1161/CIR.0000000000001078

[3] Yusuf, S., et al. (2004). Effect of potentially modifiable risk factors associated with myocardial infarction in 52 countries (the INTERHEART study): Case-control study. The Lancet, 364(9438), 937–952. https://doi.org/10.1016/S0140-6736(04)17018-9

[4] Lichtenstein, A. H., et al. (2021). 2021 Dietary Guidance to Improve Cardiovascular Health: A Scientific Statement From the American Heart Association. Circulation. https://doi.org/10.1161/CIR.0000000000001031

[5] Bull, F. C., et al. (2020). World Health Organization 2020 guidelines on physical activity and sedentary behaviour. British Journal of Sports Medicine, 54, 1451–1462. https://doi.org/10.1136/bjsports-2020-102955